

Homework 1. PHY 564 August 31 2015

Due September 9, 2015

Problem 1. 2 points. Lorentz transformations

Show explicitly that two successive Lorentz transformations in the same direction are equivalent to a single Lorentz transformation with $v = \frac{v_1 + v_2}{1 + (v_1 v_2 / c^2)}$.

Problem 2. 2 points. 4-invariants

Show that trace of a tensor is 4-invariant, i.e. $F_i^i \equiv \sum_{i=0}^3 F_i^i = inv$.

Problem 3. Lorentz group

a) **5 points.** For the Lorentz boost and rotation matrices \mathbf{K} and \mathbf{S} show that

$$\begin{aligned} (\vec{\epsilon} \vec{\mathbf{S}})^3 &= -\vec{\epsilon} \vec{\mathbf{S}}; (\vec{\epsilon} \vec{\mathbf{K}})^3 = \vec{\epsilon} \vec{\mathbf{K}}; \forall \vec{\epsilon} = \vec{\epsilon}^*; |\vec{\epsilon}| = 1; \\ \text{or } (\vec{a} \vec{\mathbf{S}})^3 &= -\vec{a} \vec{\mathbf{S}} \cdot \vec{a}^2; (\vec{a} \vec{\mathbf{K}})^3 = \vec{a} \vec{\mathbf{K}} \cdot \vec{a}^2; \forall \vec{a} = \vec{a}. \end{aligned}$$

b) **5 points.** use this results to show that

$$\begin{aligned} e^{\vec{\omega} \vec{\mathbf{S}}} &= I + \frac{\vec{\omega} \vec{\mathbf{S}}}{|\vec{\omega}|} \sin|\vec{\omega}| + \frac{(\vec{\omega} \vec{\mathbf{S}})^2}{\vec{\omega}^2} (\cos|\vec{\omega}| - 1); \\ e^{\vec{\beta} \vec{\mathbf{K}}} &= I + \frac{\vec{\beta} \vec{\mathbf{K}}}{|\vec{\beta}|} \sinh|\vec{\beta}| + \frac{(\vec{\beta} \vec{\mathbf{K}})^2}{\vec{\beta}^2} (\cosh|\vec{\beta}| - 1); \end{aligned}$$

Draw connection to Lorentz transformations (e.g. boosts and rotations).

Note: in original problem there was typo “-“ instead of “+” before *sin* and *sinh*. Everybody who found correct sign had extra 10 points!

With solutions:

Problem 1. Show explicitly that two successive Lorentz transformations in the same direction are equivalent to a single Lorentz transformation with $v = \frac{v_1 + v_2}{1 + (v_1 v_2 / c^2)}$.

Solution: Each Lorentz transformations along x-axis corresponds to the block-diagonal matrix with parameterization of :

$$L_i = \begin{bmatrix} L_i & O \\ O & I \end{bmatrix}; L_i = \gamma_i \begin{bmatrix} 1 & \beta_i \\ \beta_i & 1 \end{bmatrix}; O = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}; I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}; \det L_i = \gamma_i^2 (1 - \beta_i^2) = 1$$

and we should find parameters of L by brining it to the same form

$$L = \begin{bmatrix} L & O \\ O & I \end{bmatrix} = L_2 L_1 = \begin{bmatrix} L_2 & O \\ O & I \end{bmatrix} \begin{bmatrix} L_1 & O \\ O & I \end{bmatrix} = \begin{bmatrix} L_2 L_1 & O \\ O & I \end{bmatrix}; L = L_2 L_1.$$

The fact that $\det L = \gamma^2 (1 - \beta^2) = 1$ for any L is taking care of the rest:

.#

$$L = L_2 L_1 = \gamma_1 \gamma_2 \begin{bmatrix} 1 & \beta_2 \\ \beta_2 & 1 \end{bmatrix} \begin{bmatrix} 1 & \beta_1 \\ \beta_1 & 1 \end{bmatrix} = \gamma_1 \gamma_2 \begin{bmatrix} 1 + \beta_1 \beta_2 & \beta_1 + \beta_2 \\ \beta_1 + \beta_2 & 1 + \beta_1 \beta_2 \end{bmatrix} = \gamma_1 \gamma_2 (1 + \beta_1 \beta_2) \begin{bmatrix} 1 & \frac{\beta_1 + \beta_2}{1 + \beta_1 \beta_2} \\ \frac{\beta_1 + \beta_2}{1 + \beta_1 \beta_2} & 1 \end{bmatrix}$$

Problem 2. Show that trace of a tensor is 4-invariant, i.e. $F^i_i \equiv \sum_i F^i_i = \text{inv.}$

Solution:
$$\text{Trace}(F') = F'^i_i = \frac{\partial x'^i}{\partial x^k} \frac{\partial x^j}{\partial x'^i} F^k_j = \frac{\partial x^j}{\partial x^k} F^k_j = \delta^j_k F^k_j = F^k_k = \text{Trace}(F) \#$$

Problem 3. Lorentz group

c) **5 points.** For the Lorentz boost and rotation matrices \mathbf{K} and \mathbf{S} show that

$$\begin{aligned} (\vec{\varepsilon}\vec{\mathbf{S}})^3 &= -\vec{\varepsilon}\vec{\mathbf{S}}; (\vec{\varepsilon}\vec{\mathbf{K}})^3 = \vec{\varepsilon}\vec{\mathbf{K}}; \forall \vec{\varepsilon} = \vec{\varepsilon}^*; |\vec{\varepsilon}| = 1; \\ \text{or } (\vec{a}\vec{\mathbf{S}})^3 &= -\vec{a}\vec{\mathbf{S}} \cdot \vec{a}^2; (\vec{a}\vec{\mathbf{K}})^3 = \vec{a}\vec{\mathbf{K}} \cdot \vec{a}^2; \forall \vec{a} = \vec{a}. \end{aligned}$$

d) **5 points.** use this results to show that

$$\begin{aligned} e^{\vec{\omega}\vec{\mathbf{S}}} &= I + \frac{\vec{\omega}\vec{\mathbf{S}}}{|\vec{\omega}|} \sin|\vec{\omega}| + \frac{(\vec{\omega}\vec{\mathbf{S}})^2}{\vec{\omega}^2} (\cos|\vec{\omega}| - 1); \\ e^{\vec{\beta}\vec{\mathbf{K}}} &= I + \frac{\vec{\beta}\vec{\mathbf{K}}}{|\vec{\beta}|} \sinh|\vec{\beta}| + \frac{(\vec{\beta}\vec{\mathbf{K}})^2}{\vec{\beta}^2} (\cosh|\vec{\beta}| - 1); \end{aligned}$$

Draw connection to Lorentz transformations (e.g. boosts and rotations).

Solution: it is possible to do it by multiplying three matrices and getting confirmation. Otherwise, we can test that:

$$\begin{aligned} (\vec{a}\vec{\mathbf{K}})^3 &= (\vec{a}\vec{\mathbf{K}})^2 \cdot \vec{a}\vec{\mathbf{K}}; \\ K_\alpha K_\beta &= \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \delta_{\alpha\beta} + \begin{bmatrix} 0 & 0 \\ 0 & u_{\chi\varepsilon} \end{bmatrix} \delta_{\chi\alpha} \delta_{\varepsilon\beta}; u = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix} \end{aligned}$$

and use it to calculate square of the matrix:

$$\begin{aligned} (\vec{a}\vec{\mathbf{K}})^2 &\equiv \sum_{\alpha, \beta=1,2,3} a_\alpha a_\beta K_\alpha K_\beta = \begin{bmatrix} \vec{a}^2 & 0 \\ 0 & 0 \end{bmatrix} + \sum_{\alpha, \beta=1,2,3} \begin{bmatrix} 0 & 0 \\ 0 & a_\alpha a_\beta u_{\chi\varepsilon} \end{bmatrix} \delta_{\chi\alpha} \delta_{\varepsilon\beta} = \vec{a}^2 I + X; \\ X &= \left(\sum_{\alpha, \beta=1,2,3} \begin{bmatrix} 0 & 0 \\ 0 & a_\alpha a_\beta u_{\chi\varepsilon} \end{bmatrix} \delta_{\chi\alpha} \delta_{\varepsilon\beta} - \vec{a}^2 \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} \right). \end{aligned}$$

First term gives us desirable answer if product of matrix X and $\vec{a}\vec{\mathbf{K}}$ is zero. It is easy to show:

$$\begin{aligned} \vec{a}\vec{\mathbf{K}} &= \begin{bmatrix} 0 & \vec{a} \\ \vec{a} & 0_{3 \times 3} \end{bmatrix} - \vec{a}^2 \begin{bmatrix} 0 & 0 \\ 0 & I \end{bmatrix} \cdot \begin{bmatrix} 0 & \vec{a} \\ \vec{a} & 0_{3 \times 3} \end{bmatrix} = -\vec{a}^2 \begin{bmatrix} 0 & 0 \\ \vec{a} & 0_{3 \times 3} \end{bmatrix}; \\ \sum_{\alpha, \beta=1,2,3} a_\alpha a_\beta \delta_{\chi\alpha} \delta_{\varepsilon\beta} \begin{bmatrix} 0 & 0 \\ 0 & u_{\chi\varepsilon} \end{bmatrix} \cdot \begin{bmatrix} 0 & \vec{a} \\ \vec{a} & 0_{3 \times 3} \end{bmatrix} &= \begin{bmatrix} 0 & 0 \\ \vec{b} & 0_{3 \times 3} \end{bmatrix} \quad \cdot \#K \\ b_\chi &= \sum_{\alpha, \beta, \varepsilon=1,2,3} a_\alpha a_\beta \delta_{\chi\alpha} \delta_{\varepsilon\beta} a_\varepsilon = a_\chi \sum_{\alpha, \beta, \varepsilon=1,2,3} a_\beta \delta_{\varepsilon\beta} a_\varepsilon = a_\chi \cdot \vec{a}^2 \Rightarrow \vec{b} = \vec{a} \cdot \vec{a}^2 \end{aligned}$$

For \mathbf{S} it is even easier, noting that it is already block-diagonal matrix:

$$S_\alpha = e_{\alpha\beta\gamma} \begin{bmatrix} 0 & 0 \\ 0 & u_{\beta\gamma} \end{bmatrix}$$

and further we can drop all time components operating with 3x3 matrix:

$$[S_\alpha]_{\beta\gamma} = e_{\alpha\beta\gamma}; [\vec{a}\vec{S}]_{\beta\gamma} = a_\alpha e_{\alpha\beta\gamma};$$

$$(\vec{a}\vec{S})^2_{\beta\eta} = [a_\alpha a_\epsilon S_\alpha S_\epsilon] = a_\alpha a_\epsilon e_{\alpha\beta\gamma} e_{\epsilon\eta\gamma}; e_{\alpha\beta\gamma} e_{\epsilon\eta\gamma} = -e_{\alpha\beta\gamma} e_{\eta\epsilon\gamma} = -\delta_{\alpha\epsilon} \delta_{\beta\eta} + \delta_{\alpha\eta} \delta_{\beta\epsilon};$$

$$\delta_{\alpha\epsilon} a_\alpha a_\epsilon = \vec{a}^2; a_\alpha a_\epsilon \delta_{\alpha\eta} \delta_{\beta\epsilon} = a_\eta a_\beta; (\vec{a}\vec{S})^2_{\beta\eta} = I\vec{a}^2 + a_\beta a_\eta; a_\beta a_\eta a_\mu e_{\mu\eta\theta} \equiv 0!$$

which is equivalent to

$$(\vec{a}\vec{S})^2 (\vec{a}\vec{S}) = -\vec{a}^2 (\vec{a}\vec{S}) \quad \#S$$

b) is trivial for any matrix

$$M^3 = (-1)^n x^2 M; n = 0, 1$$

which also means that

$$M^4 = (-1)^n x^2 M^2;$$

Separating series into zero order, odd and even terms:

$$e^M = \sum_{k=0}^{\infty} \frac{M^k}{k!} = I + \sum_{k=0}^{\infty} \frac{M^{2k+1}}{(2k+1)!} + \sum_{k=1}^{\infty} \frac{M^{2k}}{(2k)!}$$

and then use induction principle to remove all powers higher than two:

$$\sum_{k=0}^{\infty} \frac{M^{2k+1}}{(2k+1)!} = \sum_{k=0}^{\infty} \frac{(M^2)^k}{(2k+1)!} M = M \sum_{k=0}^{\infty} \frac{\{(-1)^n x^2\}^k}{(2k+1)!};$$

$$\sum_{k=1}^{\infty} \frac{M^{2k}}{(2k)!} = \sum_{k=1}^{\infty} \frac{(M^2)^k}{(2k)!} = M^2 \sum_{k=1}^{\infty} \frac{\{(-1)^n x^2\}^k}{(2k)!}$$

brining the rest of the problem to known exponents:

$$i^n x \sum_{k=0}^{\infty} \frac{\{(-1)^n x^2\}^k}{(2k+1)!} = \frac{1}{2} \{e^{i^n x} - e^{-i^n x}\};$$

$$1 + (-1)^n x^2 \sum_{k=1}^{\infty} \frac{\{(-1)^n x^2\}^k}{(2k)!} = \frac{1}{2} \{e^{i^n x} + e^{-i^n x}\}$$

Therefore, both cases are identical with exception of the split between regular sin/cos and their hyperbolic twins.

In addition:

$$M = \vec{a}\vec{S} \Rightarrow x = |\vec{a}|; \Rightarrow \frac{M}{x} = \frac{\vec{a}\vec{S}}{|\vec{a}|} = \hat{e}\vec{S};$$

$$M = \vec{a}\vec{K} \Rightarrow x = |\vec{a}|; \Rightarrow \frac{M}{x} = \frac{\vec{a}\vec{K}}{|\vec{a}|} = \hat{e}\vec{K}; \#\#$$

$$\hat{e} \rightarrow \vec{\beta}; |\vec{a}| \rightarrow \zeta$$

What is left? Question about general expression for

$$e^{\vec{a}\vec{S}+\vec{b}\vec{K}} = \sum_{n=0}^{\infty} \frac{(\vec{a}\vec{S}+\vec{b}\vec{K})^n}{n!}.$$