

Homework 4

Prelude: Many elements of accelerators are straight – e.g. coordinate system is simply Cartesian ($x, y, s=z$). It allows you to forget about curvilinear coordinates and use simple div and curl and Laplacian... Many of them are DC - e.g. either with constant or nearly constant EM fields. Again, Maxwell equations without time derivatives – EM static. Furthermore, many of them are also long – e.g. have a constant cross-section with transverse size much smaller than the length of the element. It means that you can drop derivatives over z . Finally, all current and charges generating field are outside of the vacuum where particles propagate – e.g. Maxwell static equations are also homogeneous – charge and current densities are zero! It should come as no surprise – everybody like to have a solvable problem to rely upon.

Static electric and magnetic fields in vacuum can be described as gradients of a scalar potential:

$$\vec{E} = \vec{\nabla} \phi_E; \quad \vec{B} = \vec{\nabla} \phi_M.$$

While this is well-known for static electric field, it is less known for a static magnetic field in vacuum! – it is result of

Since $\vec{\nabla} \cdot \vec{B} = 0$ and in vacuum $\vec{\nabla} \cdot \vec{E} = 0$, we got in Cartesian coordinates systems

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2} \right) \phi_{E,M} = 0$$

Problem 1. 5 points. Long elements.

(a) use electro-static equations for a long uniform electric element and show that

$$\vec{E} = \vec{\nabla} \operatorname{Re} \left[a_n (x + iy)^n \right] \quad (1)$$

satisfy static Maxwell equations with a_n being a complex number. Electric elements with real a_n call regular elements (they have plane symmetry!), element with imaginary a_n are called skew .

(b) use magneto-static equations for a long uniform magnetic element

$$\vec{B} = \vec{\nabla} \operatorname{Re} \left[b_n (x + iy)^n \right] \quad (2)$$

satisfy static Maxwell equations with b being a complex number. Magnetic elements with imaginary b_n call regular elements (they have plane symmetry!), element with real b_n are called skew.

(c) show that arbitrary combination of elements from (1) and (2) is also a solution of electrostatic equations.

Hint: do not forget to prove $\vec{\nabla} \cdot \vec{E} = 0$; $\vec{\nabla} \times \vec{E} = 0$; $\vec{\nabla} \cdot \vec{B} = 0$; $\vec{\nabla} \times \vec{B} = 0$.

Note: elements with various n have specific names: $n=1$ – dipole, $n=2$ – quadrupole, $n=3$ – sextupole, $n=4$ – octupole, Or $2n$ -pole element. Term “skew” is added as needed to names of quadrupole and higher order element. It also obvious that an arbitrary $2n$ -pole “element” can be constricted as combination a regular and a skew fields.

Problem 2. 10 points. Edge effects.

- (a) **5 points.** Continue with Cartesian (x,y,s=z) coordinates for a straight element. But assume now that field in this element depends on z;

$$\vec{E}, \vec{B} = \vec{\nabla} \operatorname{Re} \left[a_n(z) (x + iy)^n \right] \quad (3)$$

Show that such elements will generate terms in the field which are not a higher order multipoles (1) or (2). Prove that a sum of higher order multi-poles with amplitudes dependent on z cannot be a solution for edge field.

- (b) **5 points.** In (a) you proved that simple combination of field multipoles cannot describe the edge of a magnet. Let expand the potential in transverse direction while keeping arbitrary dependence along the beam propagating axis (s=z)

$$\varphi = \sum_{n+m=k}^{\infty} a_{nm}(z) x^n y^m$$

and derive the condition (connections) between functions $a_{nm}(z)$ coming from $\Delta\varphi = 0$.

Problem 3. 8 points. Prove what we discussed in class:

$$\det[I + \varepsilon A] = 1 + \varepsilon \cdot \operatorname{Trace}[A] + O(\varepsilon^2)$$

where I is unit $n \times n$ matrix, A is an arbitrary $n \times n$ matrix and ε is infinitesimally small real number. Term $O(\varepsilon^2)$ means that it contains second and higher orders of ε .

Hint: first, first look on the product of diagonal elements $\prod_{m=1}^n (1 + \varepsilon a_{mm})$ in $\det[I + \varepsilon A]$ in the first order of ε . Then prove that contributions to determinant from non-diagonal terms $a_{km}; k \neq m$ is $O(\varepsilon^2)$ or higher order of ε .