

STRONG FOCUSING SYNCHROTRON

A BRIEF INTRODUCTION


- ORIGINS, PRINCIPLE
- COMBINED/SEPARATED FUNCTION
- SF-SYNCHROTRON TODAY

Bibliography

- A. Sessler, E. Wilson, Engines of Discovery, World Scientific (2007)
- M.S.~Livingston, The Development of High-Energy Accelerators, Dover Pub. Inc., NY (1966).
- CERN Accelerator School archives
- JACoW <http://www.jacow.org/>
- Joint Universities Accelerator School lectures
<http://www.esi-archamps.eu/Thematic-Schools/Discover-JUAS>
- USPAS archives
- National Lab sites, US, EU
- CERN documentation web sites
- BNL's Flickr photo gallery
- Wikipedia
- G.~Leleux, Circular accelerators, INSTN lectures, SATURNE Laboratory, CEA Saclay (Juin 1978).

Synchrotron landscape, when strong focusing was invented, 1950

Cosmotron at BNL, 1952-1968, 3.3 GeV,
the first GeV+ accelerator
(beam to target, cosmic rays' mesons,
heavy unstable particles),



occupied the front of the scene.

and Bevatron at Berkley, 1954-1993,
6 GeV, 10,000 tons of iron (discovery
of antiproton, of antineutron),



Even more ! In spite of that invention:

Synchrophasatron in Dubna (10GeV, 1957-2003!), **Saturne in France (3GeV, 1958)**, ZGS at Argonne (12GeV, 1963!-1979), **Nimrod in the UK (8 GeV, 1964!-1978)** would be built.



Genesis


- Strong focusing was patented in the early 1950s, in Greece and in the USA
- At BNL it was desired to alternate the COSMOTRON C-shaped yokes opening (all were outward), looking alternately outward and inward ... It was realized that nothing precluded strongly increasing the gradient, from its weak $0 < n < 1$ to a strong $|n| \gg 1$ with alternate sign. That's how it was discovered there in 1953
- Visitors from CERN brought BNL's idea back home, this led to the **CERN PS, 25 GeV. PS start-up: 1959.**

gamma-transition was an issue... it was solved on the fly by the PS group

Today CERN PS role still paramount, part of the injector chain to LHC
- **BNL AGS start up: 1960.**

AGS role still paramount today: RHIC injector, and part of the future EIC


A simple principle: A strong index, and alternating sign



Just like in geometrical optics:
a doublet satisfies $1/f = 1/f_1 + 1/f_2$, designed converging, here

Strong index dipole $n = \rho/B dB/dx$ + alternating gradient


Hyperbolic gap, $V = a xy$



PS or AGS, 30 GeV: few cm diameter vacuum chamber



Compare dipole size:

Cosmotron, 3 GeV:
1.22m x 0.22m vacuum chamber



FFAG SF Optics:


Sign of B alternates



Compare SATURNE 1, weak focusing and SATURNE 2, strong focusing



Mimas injector of polarized particles, of the Saturn Synchrotron at the Atomic Energy Center (CEA) in Saclay. First beam March 02, 1988 [License](#)



SATURNE 2, second (after ZGS) polarized ion synnchrotron. Same energy as SAT1: 3GeV.

***Same location,
same circumference (109
vs 105 m),
same energy (3 GeV)***






The concept evolved, from “combined function” to “separated function” optics

Dipole: steering



Quadrupole:
strong focusing




Parabolic
equipotential:

$$V=Gxy$$

$$B_x=dV/dx=Gy$$

$$B_y=dV/dy=Gx$$

Separated function optics at LEIR



Cryo-magnetism allows high field B (~ 8 T at LHC, 4 T at RHIC)



And high field gradient dB/dx (~ 250 T/m)

LHC, circumference 27km, $E=7$ TeV:


2-in-1 dipole field 8.32 T (1232 units)

Quadrupole gradient 225 T/m (392 units)


This is a cross section of a main quadrupole of the LHC at CERN: $223 \text{ T/m} \times 3.2 \text{ m}$





LHC DIPOLE : STANDARD CROSS-SECTION



**Aperture
 $\Phi 56\text{mm}$**





**The future of HEP:
still strong-focusing optics,
still similar size magnet aperture (cms) and beam (mm)**

FCC-ee

