1. The energy loss per turn is given by

$$U_0 = \frac{e^2 \beta^3 \gamma^4}{3\varepsilon_0 \rho} . \tag{1}$$

With $\rho=2.22m$ and $\gamma=1GeV\,/\,0.511MeV=1957$, eq. (1) yields

$$U_0 = \frac{e^2 \beta^3 \gamma^4}{3\varepsilon_0 \rho} = 39.6 KeV = 6.33 \times 10^{-15} J \quad .$$
 (2)

The critical photon energy is given by

$$E_c = \hbar \omega_c , \qquad (3)$$

where \hbar is the denoted Planck constant and

$$\omega_c = \frac{3}{2} \gamma^3 \frac{c}{\rho} \approx 1.512 \times 10^{18} \, rad \, / \, s \tag{4}$$

is the critical angular frequency of the synchrotron radiation. Inserting eq. (4) into eq. (3) yields

$$E_c \approx 0.996 KeV = 1.594 \times 10^{-16} J$$
 (5)

The total synchrotron radiation power for a beam is given by the 1-turn energy loss of all particles in the ring divided by the time it takes for one circulation (i.e. the revolution period)

$$P_{beam} = \left(U_0 \cdot N_{ring}\right) \frac{1}{T_{rev}} = \left(U_0 \cdot \frac{I_b}{e} T_{rev}\right) \frac{1}{T_{rev}} = U_0 \frac{I_b}{e} .$$
(6)

where $N_{ring} = I_b T_{rev} / e$ is the total number of electrons in the ring. Inserting eq. (2) and $I_b = 200 mA$ into eq. (6) give

$$P_{beam} \approx 7.91 KW \,. \tag{7}$$



Since the two intersection points are on the light-cone opened-up by $x = (x_0, \vec{x})$, they satisfy the following equation:

$$(x_0 - X_0) - \sqrt{(X_1 - x_1)^2 + (X_2 - x_3)^2 + (X_3 - x_3)^2} = 0,$$
(8)

and

$$(x_0 - Y_0) - \sqrt{(Y_1 - x_1)^2 + (Y_2 - x_3)^2 + (Y_3 - x_3)^2} = 0.$$
(9)

Subtracting eq. (9) with eq. (8) yields

$$Y_0 - X_0 = \sqrt{(X_1 - x_1)^2 + (X_2 - x_2)^2 + (X_3 - x_3)^2} - \sqrt{(Y_1 - x_1)^2 + (Y_2 - x_2)^2 + (Y_3 - x_3)^2} .$$
(10)

The three points \vec{x} , \vec{X} and \vec{Y} form a triangle and since the difference in the length of any two sides of a triangle is always smaller than the length of the third side, it follows from eq. (10)

$$Y_0 - X_0 \le \sqrt{\left(X_1 - Y_1\right)^2 + \left(X_2 - Y_2\right)^2 + \left(X_3 - Y_3\right)^2} \quad . \tag{11}$$

The time it takes for the particle to get from \vec{X} to \vec{Y} is given by

$$\Delta t = \frac{Y_0 - X_0}{c} , \qquad (12)$$

and hence the average velocity of the particle during its travelling from \vec{X} to \vec{Y} is

$$\left\langle v_{particle} \right\rangle = \frac{\sqrt{\left(X_{1} - Y_{1}\right)^{2} + \left(X_{2} - Y_{2}\right)^{2} + \left(X_{3} - Y_{3}\right)^{2}}}{\Delta t} = \frac{c\sqrt{\left(X_{1} - Y_{1}\right)^{2} + \left(X_{2} - Y_{2}\right)^{2} + \left(X_{3} - Y_{3}\right)^{2}}}{Y_{0} - X_{0}} .$$
(13)

According to eq. (11), the following relation holds

$$\frac{\sqrt{\left(X_{1}-Y_{1}\right)^{2}+\left(X_{2}-Y_{2}\right)^{2}+\left(X_{3}-Y_{3}\right)^{2}}}{Y_{0}-X_{0}} \ge 1 , \qquad (14)$$

and inserting eq. (14) into eq. (13) yields

$$\left\langle v_{particle} \right\rangle = c \frac{\sqrt{\left(X_1 - Y_1\right)^2 + \left(X_2 - Y_2\right)^2 + \left(X_3 - Y_3\right)^2}}{Y_0 - X_0} \ge c$$
 (15)

Eq. (15) violates special relativity and hence the trajectory of a particle cannot intersect a light-cone twice.

3. The angular distribution of radiation power is given by

$$\frac{dP(t_r)}{d\Omega} = \frac{1}{4\pi\varepsilon_0} \frac{e^2}{4\pi c} \frac{\dot{\beta}^2}{\left(1 - \beta\cos\theta\right)^3} \left[1 - \frac{\sin^2\theta\cos^2\phi}{\gamma^2\left(1 - \beta\cos\theta\right)^2} \right].$$
 (1)

For $\frac{1}{\gamma^{^4}} <\!\!< \!\theta <\!\!< \!1$ and $\gamma \!>\!\!> \!1$, we can use the following approximation

$$1 - \beta \cos \theta \approx 1 - \beta \left(1 - \frac{1}{2} \theta^2 \right)$$

$$= 1 - \beta + \frac{1}{2} \beta \theta^2$$

$$= \frac{1}{\gamma^2 (1 + \beta)} + \frac{1}{2} \theta^2$$

$$= \frac{1}{\gamma^2} \left[\frac{1}{2 - (1 - \beta)} \right] + \frac{1}{2} \theta^2 , \qquad (2)$$

$$\approx \frac{1}{2\gamma^2} \left[1 + \frac{1 - \beta}{2} \right] + \frac{1}{2} \theta^2$$

$$\approx \frac{1}{2\gamma^2} \left[1 + \frac{1}{4\gamma^2} + \dots \right] + \frac{1}{2} \theta^2$$

$$\approx \frac{1}{2\gamma^2} + \frac{1}{2} \theta^2$$

and eq. (1) becomes

$$\frac{dP(t_r)}{d\Omega} \approx \frac{1}{4\pi\varepsilon_0} \frac{2e^2}{\pi c} \frac{\gamma^6 \dot{\beta}^2}{\left(1+\gamma^2 \theta^2\right)^3} \left[1 - \frac{4\gamma^2 \theta^2 \cos^2 \phi}{\left(1+\gamma^2 \theta^2\right)^2}\right].$$
 (3)

Since the factor inside the square bracket is between 0 and 1, the angular width of eq. (3) is determined by the factor $(1 + \gamma^2 \theta^2)^{-3}$, i.e. the radiation power drops substantially when $\theta \ge \frac{1}{\gamma}$.